arXiv:hep-th/0605233v3 2 Jul 2006

Remarks on four dimensional Euclidean gravity without Wick rotation

Jens Köplinger*
406 Aberdeen Ter, Greensboro, NC 27403, USA

(Dated: 20 May 2006)

In reference to S. W. Hawking's article "Information Loss in Black Holes" [S. W. Hawking, Phys. Rev. D 72 (2005) 084013], where a four dimensional Euclidean spacetime without Wick rotation is adopted for quantum gravity, an arithmetic with multiplicative modulus is mentioned here which incorporates both a hyperbolic (Minkowski) and circular (Euclidean) metric: The 16 dimensional conic sedenion number system is built on nonreal square roots of +1 and -1, and describes the Dirac equation through its 8 dimensional hyperbolic octonion subalgebra [J. Köplinger, Appl. Math. Comput. (2006), in print, doi: 10.1016/j.amc.2006.04.005]. The corresponding circular octonion subalgebra exhibits Euclidean metric, and its applicability in physics is being proposed for validation. In addition to anti-de Sitter (AdS) spacetimes suggested by Hawking, these conic sedenions are offered as computational tool to potentially aid a description of quantum gravity on genuine four-dimensional Euclidean spacetime (without Wick rotation of the time element), while being consistent with canonical spacetime metrics.

PACS numbers: 04.70.Dy
Keywords: black holes; space-time configurations; topology; quantum gravity; hypernumbers; conic complex numbers; sedenions

In [1] it is shown that "the path integral over topologically trivial metrics like periodically identified AdS is unitary". Such topologically trivial metrics allow to describe wave functions as functional of the geometries of space-like surfaces on slices with constant time. With evidence that all other, non-trivial black hole topologies must lead to decay of correlation functions, it is concluded for an observation at infinity that the whole path integral will be unitary, and information that enters a black hole may return to our universe.

In executing these deductions, gravity is treated on a four dimensional Euclidean metric, but without Wick rotation of the time interval as one may expect. Anti-de Sitter coordinate patches reduce to Minkowski spacetime at infinity while also incorporating four dimensional Euclidean metric otherwise. They therefore satisfy mathematical description of above deduction, which "seems to be well formulated only in asymptotically AdS spacetimes".

It is now mentioned that use of the non-traditional number system of conic sedenions (e.g. [2, 3]) may pose an additional option for formulation of multiple trivial geometries on the same physical universe, in particular Minkowski and Euclidean spacetime, with certain superpositions thereof.

Conic sedenions contain octonion subalgebras with hyperbolic (Minkowski) and circular (Euclidean) geometries[6]. They form a 16 dimensional arithmetic which has a multiplicative modulus, but is non-commutative and non-associative. They cannot be obtained through matrix-extension of traditional complex numbers or Cayley-Dickson constructs like traditional quaternions, octonions, and further 2^n dimensional number systems on 1 real and $(2^n - 1)$ imaginary bases (roots of -1). Conic sedenions, instead, are built on 1 real basis, 7 imaginary bases i_n (with $i_n^2 = -1$), 7 "counterimaginary" [2] bases ε_n (nonreal $\sqrt{+1}$, $\varepsilon_n^2 = 1$), and 1 compound basis i_0 (with $i_0 := i_1\varepsilon_1 = i_2\varepsilon_2 = ... = i_7\varepsilon_7$).

It is shown in [5] that while the hyperbolic octonion subalgebra of conic sedenions can represent the classical ("hyperbolic") Dirac equation without field in the simple form

$$\nabla_{\text{hyp8}}\psi_{\text{hyp8}} = 0 \tag{1}$$

, rotation in the $(1, i_0)$ plane by a real angle α transforms (1) into a counterpart on circular (Euclidean) octonion algebra. Conic sedenions are concluded there as the unifying number system which encompasses both geometries, and the rotation $\exp i_0\alpha$ is proposed as "symmetry transformation of a new class formed by genuine conic complex rotations".

Modularity of conic sedenions should warrant unambiguous description of a quantum system, and accessibility through measurement. They are therefore proposed as an arithmetic that potentially offers the benefit of describing physical law, next to AdS formulations.

---

*Electronic address: jens@prisage.com

Acknowledgments

I am thankful to Kevin Carmody for his generous help with hypernumber arithmetics. For further information on hypernumbers see e.g. http://www.kevincarmody.com/math/hypernumbers.html .

[1] S. W. Hawking, *Information Loss in Black Holes*, Phys. Rev. D 72 (2005) 084013, preprint arXiv:hep-th/0507171

[2] K. Carmody, *Circular and Hyperbolic Quaternions, Octonions, and Sedenions*, Appl. Math. Comput. 28 (1988) 47-72, preprint http://www.kevincarmody.com/math/hypernumbers.html

[3] K. Carmody, *Circular and Hyperbolic Quaternions, Octonions, and Sedenions - Further Results*, Appl. Math. Comput. 84 (1997) 27-47, preprint http://www.kevincarmody.com/math/hypernumbers.html

[4] M. Gogberashvili, *Octonionic Electrodynamics*, J. Phys. A: Math. Gen. 39 (2006) 7099-7104, preprint arXiv:hep-th/0512258

[5] J. Köplinger, *Dirac Equation on Hyperbolic Octonions*, Appl. Math. Comput. (2006), in print, doi: 10.1016/j.amc.2006.04.005, preprint http://www.prisage.com/P

[6] Hyperbolic octonions are computationally identical to split-octonions (e.g. [4]). In addition, conic sedenions contain conic octonion subalgebras which exhibit a mixed geometry.